

**Central Bureau for Astronomical Telegrams**  
**INTERNATIONAL ASTRONOMICAL UNION**

Mailstop 18, Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.  
 IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions)  
 CBAT@CFA.HARVARD.EDU (science)  
 URL <http://cfa-www.harvard.edu/iau/cbat.html> ISSN 0081-0304  
 Phone 617-495-7440/7244/7444 (for emergency use only)

*COMET 9P/TEMPEL*

N. Biver, D. Bockelée-Morvan, P. Colom, J. Crovisier, and A. Lecauchaux, Observatoire de Paris; and G. Paubert, Institut de Radioastronomie Millimétrique (IRAM), report: “Comet 9P/Tempel was observed with the IRAM 30-m radio telescope from May 4.8 to 9.0 UT. The HCN  $J = 3-2$  and  $J = 1-0$  lines were detected with mean integrated intensities of  $111 \pm 8$  and  $31 \pm 4$  mK km/s, respectively. Day-to-day variations in the HCN line intensities are observed and reveal periodic variability in HCN production from 5 to  $10 \times 10^{24}$  molecules/s. The periodicity is 1.7 days, consistent with the estimated nucleus rotational period. H<sub>2</sub>S and CH<sub>3</sub>OH were also marginally detected; CO, CS and H<sub>2</sub>CO were searched for and undetected. The production-rate ratios, or their upper limits, of these molecules relative to HCN are 5.5, 21, < 110, < 1.1, and < 17, respectively. All lines are blue-shifted, suggesting preferential sunward outgassing.”

Total-magnitude and coma-diameter estimates made with 25×100 binoculars: Apr. 28.97 UT, 10.9, 3' (J. J. Gonzalez, Asturias, Spain); May 5.54, 10.3, – (D. A. J. Seargent, The Entrance, N.S.W.); 26.90, 10.1, 4' (M. Lehký, Hradec Králové, Czech Rep.); June 5.97, 10.1, 3'5 (Gonzalez).

*SUPERNOVAE 2005br, 2005bs, 2005cb*

M. Turatto, S. Benetti, and A. Harutyunyan, Istituto Nazionale di Astrofisica (INAF), Padova; M. Riello, Cambridge Astronomical Survey Unit, Institute of Astronomy, University of Cambridge; E. Cappellaro, INAF, Napoli; M. T. Botticella, INAF, Teramo; E. Mason, European Southern Observatory (ESO), write that that obtained spectroscopy of three southern supernovae with the ANTU Very Large Telescope (+ Fors2) at ESO on May 25.2 UT under poor weather conditions. Based on the reduced spectra (range 380–920 nm, resolution 1.0 nm), SN 2005br in IC 5084 (*IAUC* 8516) appears to be a type-Ib SN  $\sim 40$  days past maximum; the spectrum is very reddened and shows a strong interstellar Na I D absorption with EW  $\sim 0.25$  nm. SN 2005bs (*IAUC* 8517) is a normal type-Ia supernova about 30 days past maximum. SN 2005cb in NGC 6753 (*IAUC* 8530) appears to be a type-Ib/c supernova, similar to SN 1997X (Munari *et al.* 1998, *A.Ap.* **333**, 159) at about 10 days after maximum.

T. Davis, G. Anderson, and B. Schmidt, Australian National University (ANU), report that a spectrogram (range 340–920 nm) of SN 2005cb (cf. *IAUC* 8530), obtained on May 19.76 UT with the ANU 2.3-m telescope (+ DBS), reveals it to be a type-Ic supernova  $\approx 1$  week past maximum light.