

**Central Bureau for Astronomical Telegrams
INTERNATIONAL ASTRONOMICAL UNION**

Mailstop 18, Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.
IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions)
CBAT@CFA.HARVARD.EDU (science)
URL <http://www.cfa.harvard.edu/iau/cbat.html> ISSN 0081-0304
Phone 617-495-7440/7244/7444 (for emergency use only)

COMET C/2007 Y2 (McNAUGHT)

R. H. McNaught reports his discovery of a comet (as tabulated below) on CCD images taken with the 0.5-m Uppsala Schmidt telescope, where the object appears marginally, but consistently, diffuse with a coma diameter of $\sim 10''$ in good seeing. Three stacked 90-s *R* images taken by D. M. Burton, Australian National University, with the ANU 1.0-m *f*/8 reflector at Siding Spring on 2008 Jan. 1.69 show a slightly diffuse comet (slightly involved with a star) with a $3''$ tail in p.a. 20° . Following posting on the Minor Planet Center's 'NEOCP' webpage, C. Jacques and E. Pimentel write that their astrometric CCD images from Jan. 1.14 (taken with a 0.30-m *f*/3 reflector at Belo Horizonte, Brazil) show a central condensation of size $10''$.

2007	UT	α_{2000}	δ_{2000}	Mag.
Dec.	31.66576	$6^{\text{h}}45^{\text{m}}01^{\text{s}}.58$	$-50^\circ45'51''.5$	19.0

The available astrometry, the following preliminary parabolic orbital elements, and an ephemeris appear on *MPEC* 2008-A6.

$$\begin{array}{rcl}
 T = 2008 \text{ May } 18.189 \text{ TT} & \omega = 264.435 & \\
 & \Omega = 303.854 & \left. \vphantom{\begin{array}{l} \omega \\ \Omega \end{array}} \right\} 2000.0 \\
 q = 4.13214 \text{ AU} & i = 98.702 &
 \end{array}$$

SUPERNOVA 2007uy IN NGC 2770

S. Nakano, Sumoto, Japan, reports the discovery of an apparent supernova (mag 17.2) by Yoji Hirose (Tsutsumi, Chigasaki, Kanagawa-ken, Japan) on ten unfiltered CCD frames taken around 2007 Dec. 31.669 UT using a 0.35-m *f*/6.8 Schmidt-Cassegrain telescope (limiting mag 18.0); Nakano has measured mag 17.0 and the following position for the new object from a jpeg image supplied by Hirose: $\alpha = 9^{\text{h}}09^{\text{m}}35^{\text{s}}.40$, $\delta = +33^\circ07'09''.9$ (equinox 2000.0), which is $20''.6$ east and $15''.5$ south of the center of NGC 2770. Nothing is visible at this position on red Digitized Sky Survey images from 1998 and 2000, though there are apparently numerous H II regions nearby. Additional unfiltered CCD magnitudes for 2007uy: 2007 Mar. 7 and Dec. 18, [18 (Hirose); Dec. 31.786, 17.3 (K. Kadota, Ageo, Japan, 0.25-m *f*/5 reflector; position end figures $35^{\text{s}}27$, $08''9$); 2008 Jan. 1.595, 17.0 (Hirose); 1.651, 16.9 (K. Itagaki, Takanezawa station, Tochigi-ken, Japan, 0.30-reflector; position end figures $35^{\text{s}}35$, $08''9$); 1.885, 16.8 (P. Corelli, Pagnacco, Italy, 0.45-m *f*/4.5 reflector; position end figures $35^{\text{s}}17$, $08''9$).